Asthma Treatment Plan

CITY OF AUGUSTA CHILDCARE BUREAU

(Please Print)

Name	Date of Birth	Effective Date
Doctor	Parent/Guardian (if applicable)	Emergency Contact
Phone	Phone	Phone

HEALTHY

You have *all* of these:

- Breathing is good
- No cough or wheeze
- Sleep through the night
- Can work, exercise, and play

And/or Peak flow above _____

Take daily medicine(s). All metered dose inhalers (MDI) to be used with spacers.

MEDICINE	HOW MUCH to take and HOW OFTEN to take it
☐ Advair® 100, 250, 500	1 inhalation twice a day
☐ Advair® HFA 45, 115, 230	2 puffs MDI twice a day
☐ Asmanex® Twisthaler® 110, 220	1 - 2 inhalations a day
☐ Flovent® 44, 110, 220	2 inhalations twice a day
☐ Flovent® Diskus® 50 mcg	1 inhalation twice a day
☐ Pulmicort Flexhaler® 90, 180	1 - 2 inhalations once or twice a day
☐ Pulmicort Respules® 0.25, 0.5, 1.0	1 unit nebulized once or twice a day
☐ Qvar® 40, 80	2 inhalations twice a day
☐ Singulair 4, 5, 10 mg	1 tablet daily
☐ Symbicort® 80, 160	2 puffs MDI twice a day
☐ Other	

Remember to rinse your mouth after taking inhaled medicine.

If exercise triggers your asthma, take this medicine_____ ____minutes before exercise.

CAUTION

You have *any* of these:

- Exposure to known trigger
- Cough
- Mild wheeze
- Tight chest
- Coughing at night
- Other:_____

And/or Peak flow from_____ to_____

Continue daily medicine(s) and add fast-acting medicine(s).

MEDICINE	HOW MUCH to take and HOW OFTEN to take it
☐ Accuneb® 0.63, 1.25 mg	1 unit nebulized every 4 hours as needed
☐ Albuterol 1.25, 2.5 mg	1 unit nebulized every 4 hours as needed
☐ Albuterol ☐ Pro-Air ☐ Proventil®	2 puffs MDI every 4 hours as needed
☐ Ventolin® ☐ Maxair ☐ Xopenex®	2 puffs MDI every 4 hours as needed
☐ Xopenex® 0.31, 0.63, 1.25 mg	1 unit nebulized every 4 hours as needed
☐ Increase the dose of, or add:	

➡ **If fast-acting medicine is needed more than 2 times a week, except before exercise, then call your doctor.**

EMERGENCY



Your asthma is getting worse fast:

- Fast-acting medicine did not help within 15-20 minutes
- Breathing is hard and fast
- Nose opens wide
- Ribs show
- Trouble walking and talking
- Lips blue • Fingernails blue

And/or Peak flow below _____

Take these medicines NOW and call 911.

Asthma can be a life-threatening illness. Do not wait!

☐ Accuneb® 0.63, 1.25 mg	1 unit nebulized every 20 minutes
☐ Albuterol 1.25, 2.5 mg	1 unit nebulized every 20 minutes
☐ Albuterol ☐ Pro-Air ☐ Proventil®	2 puffs MDI every 20 minutes
☐ Ventolin® ☐ Maxair ☐ Xopenex®	2 puffs MDI every 20 minutes
☐ Xopenex® 0.31, 0.63, 1.25 mg	1 unit nebulized every 20 minutes
☐ Other	

Triggers

Check all items that trigger patient's asthma:

- ☐ Chalk dust
- ☐ Cigarette Smoke & second hand smoke
- ☐ Colds/Flu
- ☐ Dust mites, dust, stuffed animals, carpet
- ☐ Exercise
- ☐ Mold
- ☐ Ozone alert days
- ☐ Pests - rodents & cockroaches
- ☐ Pets - animal dander
- ☐ Plants, flowers, cut grass, pollen
- ☐ Strong odors, perfumes, cleaning products, scented products
- ☐ Sudden temperature change
- ☐ Wood Smoke
- ☐ Foods: _____ _____ _____
- ☐ Other: _____ _____ _____ _____ _____

This asthma treatment plan is meant to assist, not replace, the clinical decision-making required to meet individual patient needs.

FOR MINORS ONLY:

☐ This student is capable and has been instructed in the proper method of self-administering of the inhaled medications named above _____

☐ This student is not approved to self-medicate.

PARENT/GUARDIAN SIGNATURE_____